

Topological phases in two-dimensional arrays of parafermionic zero modes

M. Burrello, B. van Heck, and E. Cobanera

Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands.

(Dated: February 20, 2013)

It has recently been realized that zero modes with projective non-Abelian statistics, generalizing the notion of Majorana bound states, may exist at the interface between a superconductor and a ferromagnet along the edge of a fractional topological insulator (FTI). Here we study two dimensional architectures of these non-Abelian zero modes, whose interactions are generated by the charging and Josephson energies of the superconductors. We derive low-energy Hamiltonians for two different arrays of FTIs on the plane, revealing an interesting interplay between the real-space geometry of the system and its topological properties. On the one hand, in a geometry where the length of the FTI edges is independent on the system size, the array has a topologically ordered phase, giving rise to a qudit toric code Hamiltonian in perturbation theory. On the other hand, in a geometry where the length of the edges scales with system size, we find an exact duality to an Abelian lattice gauge theory and no topological order.

PACS numbers: 05.30.Pr, 74.81.Fa, 03.67.Lx, 11.15.Ha

I. INTRODUCTION

Systems that exhibit topological quantum order¹ have been a focus of attention in recent years. Part of the interest is due to the fact that they have been proposed as fault-tolerant quantum memories and platforms for quantum computation², the paradigmatic example being Kitaev's toric code³. The goal is to design architectures effectively governed by topologically-ordered Hamiltonians, where qubits may be stored and manipulated in a physically protected way.

Majorana zero modes, realized as superconducting midgap excitations in either one⁴ or two^{5,6} spatial dimensions, are promising building blocks for such architectures. Two unpaired Majorana fermions at the ends of a one-dimensional (1D) superconducting wire encode non-locally a qubit⁴ and, when allowed to move in a non-strictly 1D geometry, exhibit non-Abelian statistics⁷⁻⁹, allowing to implement a non-universal set of quantum gates through ordered exchanges of their positions. Interest in Majorana fermions has increased considerably in recent times, since there are now several experimentally accessible systems that may host these quasiparticles (see Refs. 10 and 11 for a review). A notable example is the edge of a two-dimensional (2D) topological insulator^{12,13}, which hosts gapless helical (i.e. counter-propagating) modes, in proximity to an *s*-wave superconductor (SC) and a ferromagnet (FM). The competition between the proximity-induced SC and FM pairing along the edge results in the presence of a Majorana fermion at each domain wall¹⁴.

Recently, an interesting extension of this model was proposed in Refs. 15–18. While the edge excitations of a 2D topological insulator are normal electrons, it is possible to consider instead edge quasiparticles with a fractional charge e/m , where m is an odd integer. Such gapless quasiparticles appear, for example, at the edge of the Laughlin fractional quantum Hall states, where they are described by a chiral Luttinger liquid theory^{19,20}. Due

to the absence of time-reversal symmetry, these are chiral excitations. Helical e/m quasiparticles would arise at the interface between two $\nu = 1/m$ quantum Hall liquids with Landé g -factors of opposite sign or, similarly, as a Kramers doublet at the edge of a fractional topological insulator²¹ (FTI). Also in this case, the proximity effect with superconductors and ferromagnets results in the presence of zero modes. Since the second-quantization operators associated with these zero modes inherit a fractional exchange phase $(2\pi)/(2m)$ from the unperturbed edge fields, we shall refer to them as \mathbb{Z}_{2m} parafermionic (PF) zero modes. They are projective non-Abelian anyons^{22,23}, with fusion rules that generalize those of Majorana fermions, affording extended computational power^{15,16,22,24,25}.

These superconducting zero modes realize a 1D model with \mathbb{Z}_{2m} symmetry studied by Fendley²⁶, which is an extension of Kitaev's Majorana chain model⁴ and hosts PF zero modes localized at the edges of the system. While the Kitaev chain is dual to the quantum Ising chain via a Jordan-Wigner transformation, the \mathbb{Z}_{2m} chain model is dual to the 1D chiral Potts (p -clock) model, with $p = 2m$.

Indeed, PFs as collective degrees of freedom are indeed well-known in statistical mechanics²⁷. They appear naturally in the study of the 2D p -state clock models²⁸⁻³⁰ and their quantum 1D counterparts³¹. In lattice systems, they arise as products of order and disorder operators defined for self-dual systems. Moreover, PFs admit a description in terms of a \mathbb{Z}_p invariant conformal field theory³² (CFT) featuring the PFs as primary fields. PF zero modes in superconducting systems are however related only to CFTs with unit central charge (see, for example, Ref. 33), arising from a bosonization description of the FTIs edge modes.

In light of this body of research, it is interesting to extend the recent works on \mathbb{Z}_{2m} PFs to 2D networks of superconductors. For Majorana zero modes, this question has already been addressed in literature³⁴⁻³⁶. Majorana lattice models can be mapped into Ising models, allow-




FIG. 1: The two different 2D architectures considered in this paper. They are composed of superconductors (SC, blue) and ferromagnets (FM, red) deposited on top of a 2D array of fractional topological insulators (FTI, grey). \mathbb{Z}_{2m} PF zero modes, marked as black stars, arise at each SC/FM interface along the edge. We consider two possible geometries: in panel (a), the FTIs extend for the whole length of the system, while in panel (b) the FTIs have fixed size. If we enlarged on the horizontal direction the system in panel (a), the number of FTIs would stay constant and the edge length would increase, while the vice versa would happen in panel (b). In the main text, we refer to the two architectures as the *stripe* and *tile* model respectively.

ing for a description of their phase diagram. They can exhibit topologically ordered phases and realize the toric code in a perturbative regime³⁵. The extension of these analyses to \mathbb{Z}_{2m} PFs may reveal new Hamiltonian realizations of fault-tolerant stabilizer codes^{37,38} for quantum bits with $2m$ states, and hence novel platforms for quantum memories generalizing the toric code³⁹. However, for \mathbb{Z}_{2m} PFs the extension to 2D lattices is less immediate than in the Majorana case, partially because of the connection to clock models, which are less well understood than Ising models.

In order to fill this gap, in this work we consider two distinct 2D architectures of \mathbb{Z}_{2m} PFs, shown in Fig. 1. The architectures are obtained from a pattern of superconductors and ferromagnets layered on top of an array of 2D FTIs. The only difference between the two models is the geometry of the underlying FTI array. In Fig. 1a) the array is formed by long stripes of FTIs extending for the whole length of the system, while in Fig. 1b) the stripes are cut in smaller pieces (or tiles) of fixed dimension. For ease of discussion, we shall refer to these two architectures as the *stripe* and *tile* model respectively.

Similarly to Refs.^{34–36}, the effective Hamiltonian of the two models is dictated by two mesoscopic phenomena:

1. the fractional Josephson effect, mediated by the tunneling of e/m quasiparticle between two different superconductors, and
2. the charging energy of the superconductors.

The fractional Josephson effect arising with \mathbb{Z}_{2m} PF zero modes has already been investigated in Refs.^{16,17}, while to our knowledge the interplay between PF zero modes and Coulomb energy was not considered in previous works.

While both the stripe and the tile architectures give rise to a square lattice of \mathbb{Z}_{2m} PF zero modes, and despite the fact that the effective Hamiltonian contains the same set of local interactions in both models, we will show that the different geometry of the FTI edges is crucial to determine their properties, which turn out to be quite distinct. Indeed, since different edge geometries yield a different set of commutation rules for the \mathbb{Z}_{2m} PF operators and different physical constraints on the Hilbert space, they can determine different topological properties.

The paper is organized as follows: in Sec. II we derive the effective Hamiltonian for the stripe and tile architectures, considering both Josephson and Coulomb energies, and explain the physical constraints and conservation laws specific to each array. In Sec. III we map the effective Hamiltonian into two different clock models, using two non-isomorphic sets of PF Jordan-Wigner transformations. We analyze the phase diagram of the two models in Sec. IV, where we show that the tile model realizes a qudit toric code Hamiltonian in perturbation theory while the stripe model is dual to a gauge theory. We conclude with an outlook in Sec. V.

II. EFFECTIVE HAMILTONIAN FOR 2D PARA-FERMIONIC ARCHITECTURES

In the two architectures in Fig. 1, each FM/SC interface along the edge of a FTI hosts a PF zero mode. Hence the stripe and tile architectures generate arrays of interacting \mathbb{Z}_{2m} PFs, which are protected by the superconducting and ferromagnetic gaps and thus determine the low-energy physics of the system. In this section we derive an effective Hamiltonian in terms of \mathbb{Z}_{2m} PF operators for a square lattice of dimensions $L_x \times L_y$ with open boundaries⁴⁰. Generalizations to other boundary conditions can be easily implemented.

Each PF zero mode is described by a second-quantization operator α satisfying the relations

$$\alpha^{2m} = 1 \quad (1)$$

$$\alpha^\dagger = \alpha^{2m-1} \quad (2)$$

We can associate to α and α^\dagger respectively the annihilation and creation of a charge e/m on the adjacent superconductor, in such a way that a Cooper pair is split in $2m$ quasiparticles¹⁵.




FIG. 2: A square lattice of \mathbb{Z}_{2m} PF zero modes of dimensions $L_x = 8$ and $L_y = 4$. To label the PFs, we follow the convention established in the main text: first we order the FTI edges with an index a , and then we order the PFs along each edge with an index j , starting from an arbitrary origin. Each PF zero mode is then denoted as $\alpha_{\{a,j\}}$, and all commutation rules between operators at different sites are fixed unambiguously.

Furthermore, these PF operators obey unconventional commutation rules. Denoting two different PF operators with generic, ordered labels μ and ν , we have

$$\alpha_\mu \alpha_\nu = e^{-i\epsilon_{\mu\nu}\pi/m} \alpha_\nu \alpha_\mu \quad (3)$$

$$\alpha_\mu^\dagger \alpha_\nu = e^{+i\epsilon_{\mu\nu}\pi/m} \alpha_\nu \alpha_\mu^\dagger \quad (4)$$

where $\epsilon_{\mu\nu} = -\epsilon_{\nu\mu} = \pm 1$ is a sign that must be fixed by convention. As we outline in Appendix A, Eqs. (1-4) can be derived from the underlying helical Luttinger liquid theory for the FTI edges. Note that for $m = 1$ the $\epsilon_{\mu\nu}$'s do not play any role and the equations (1-4) reproduce all the properties of Majorana fermions.

In the 1D case, μ and ν are integers denoting the positions of the PFs on a line. All signs are fixed by assigning an orientation to the line, so that $\epsilon_{\mu\nu} = \text{sgn}(\mu - \nu)$. In two dimensions the ordering procedure is slightly more complicated and proceeds in the following way.

1. We label each FTI edge of the system with an integer a , thus introducing an ordering of the edges. We also assign a counterclockwise orientation to each edge a .
2. Starting from an arbitrary origin and following the counterclockwise orientation, we label all ferromagnets along the edge with an integer $k = 1, \dots, M$ (similarly to what was done in Ref. 15). The number M is the total number of FMs along each FTI edge: note that $M = 4$ for the tile model, while $M = L_x$ for the stripe model (see Fig. 2).
3. We identify the SC/FM interfaces at the left and the right of each FM with an integer $j = 2k - 1$ or $j = 2k$ respectively.

The PFs α_μ, α_ν are thus labelled by a composite index $\mu = \{a, j\}$, $\nu = \{a', j'\}$ and we fix all the conventional signs as

$$\epsilon_{\mu\nu} = \text{sgn}(a - a') + \delta_{aa'} \text{sgn}(j - j') \quad (5)$$

In Fig. 2 we explicitly illustrate the procedure for labeling all the PFs of our square array, in both the tile and stripe architecture. Due to the different number and disposition of the FTIs, the PFs in the stripe model are actually distinct from (non-isomorphic to) the PFs in the tile model. The value of $\epsilon_{\mu\nu}$ may differ for pairs of PFs in the same site of the square lattice, and consequently the set of commutation relations Eqs. (3,4) is not the same for the two geometries.

In order to describe physical interactions between PFs, it is useful to introduce the operator

$$P_{\mu\nu} = e^{i\epsilon_{\mu\nu}\pi/2m} \alpha_\nu^\dagger \alpha_\mu, \quad (6)$$

defined for every given pair α_μ, α_ν . For $m = 1$, $P_{\mu\nu}$ represents the \mathbb{Z}_2 fermionic parity associated with two Majorana fermions. Here we are extending this notion to the \mathbb{Z}_{2m} symmetry of the PFs, and we shall refer to $P_{\mu\nu}$ as \mathbb{Z}_{2m} charge. From its Hermitian conjugate,

$$P_{\mu\nu}^\dagger = e^{-i\epsilon_{\mu\nu}\pi/2m} \alpha_\mu^\dagger \alpha_\nu \quad (7)$$

we see that it is a unitary operator,

$$P_{\mu\nu} P_{\mu\nu}^\dagger = P_{\mu\nu}^\dagger P_{\mu\nu} = 1 \quad (8)$$

Moreover $P_{\mu\nu}^{2m} = 1$. Thus its eigenvalues must be the $(2m)$ -th roots of unity,

$$\lambda_n = e^{in\pi/m}, \quad n = 0, 1, \dots, 2m - 1. \quad (9)$$

A pair α_μ, α_ν can be irreducibly represented on a $2m$ -dimensional Hilbert space, with a basis given by the states $|n\rangle$ such that $P_{\mu\nu}|n\rangle = \lambda_n|n\rangle$. The Hilbert space dimension of a square lattice of PF zero modes of size $L_x \times L_y$ is therefore $(2m)^{L_x \cdot L_y/2}$.

Now that we have set the basic algebraic rules, we can write down the physical ingredients of the model - Josephson and charging energy. These will form the basic local bonds used to write an effective 2D Hamiltonian for the PFs.

A. Josephson energy

Thanks to the presence of zero modes, phase-coherent tunneling of e/m quasiparticles may take place across the ferromagnetic region between adjacent superconductors along a common edge. The resulting Josephson effect is characterized by an anomalous periodicity of $4\pi m$ (in units of the superconducting flux quantum $\Phi_0 = h/2e$), essentially because the tunneling quasiparticle's charge is reduced by a factor $2m$ with respect to the charge of a Cooper pair^{14–17}. In other words, the anomalous period reflects the fact that the junction can be in $2m$ different states associated to the \mathbb{Z}_{2m} charge of two PFs located at its ends. Physically, these states are distinguished by the fractional spin of the ferromagnet inside the junction¹⁵, or equivalently by the number of the fractional quasiparticles trapped in it (modulo $2m$).

Using the notation introduced in Fig. 2, the \mathbb{Z}_{2m} charge of a junction situated on edge a can be written as

$$P_{\{a,2k-1\},\{a,2k\}} = e^{-i\pi/2m} \alpha_{\{a,2k\}}^\dagger \alpha_{\{a,2k-1\}} \quad (10)$$

It acts as a transfer operator, destroying one e/m charge inside the superconductor on one side of the junction and creating it on the other side. Such tunneling processes can be modeled by an effective Hamiltonian of the form

$$H_J = -\frac{J}{2} \left(e^{i(\delta-\pi)/2m} \alpha_{\{a,2k\}}^\dagger \alpha_{\{a,2k-1\}} + \text{H.c.} \right) \quad (11)$$

Here J is the tunneling strength and δ is the phase difference between the two superconductors. The tunneling Hamiltonian splits the states of the junction in $2m$ energy branches given by

$$E_{J,n} = -J \cos \left(\frac{\delta}{2m} + \frac{n\pi}{m} \right) \quad (12)$$

with $n = 0, \dots, 2m-1$.

B. Charging energy

Let us now consider a single superconducting island of our array, and let us denote with ϕ and $N = -2i \frac{d}{d\phi}$ the phase and number operators of this island. The presence of the PF zero modes becomes manifest through non-trivial (twisted) boundary conditions in the condensate ground state wave-function $\Psi(\phi)$ ⁴¹,

$$\Psi(\phi + 2\pi) = e^{i\pi q} \Psi(\phi). \quad (13)$$

Here q represents the charge in units of e inside the superconductor (modulo $2e$). The spectrum of the number operator depends on these twisted boundary conditions, since its twisted eigenfunctions $\chi_s(\phi) = e^{i(s+q/2)\phi}/\sqrt{2\pi}$ satisfy

$$N\chi_s = (2s+q)\chi_s, \quad s \in \mathbb{Z}. \quad (14)$$

In a conventional superconductor, $q = 0$, we have periodic boundary conditions, and N counts Cooper pairs. In the presence of Majoranas q may assume either value $\{0, 1\}$ giving periodic or anti-periodic boundary conditions depending on the fermion parity of the superconductor⁴¹. In the presence of \mathbb{Z}_{2m} PF zero modes, the possible values of q are extended to fractional values

$$q = \left\{ \frac{n}{m} \right\} = \left\{ 0, \frac{1}{m}, \frac{2}{m}, \dots, 1, \frac{m+1}{m}, \dots, \frac{2m-1}{m} \right\}. \quad (15)$$

The resulting boundary conditions are twisted with possible phases $e^{in\pi/m}$, and the spectrum of N is given by rational numbers with denominator m .

Ground states with different values of q are not anymore degenerate if the charging energy of the superconducting island,

$$H_{\text{ch}} = E_C (N - n_{\text{ind}})^2, \quad (16)$$

is taken into account. Here $E_C = e^2/2C$, C is the self-capacitance of the superconductor, and n_{ind} the charge (in units of e) induced on the island by nearby voltage gates.

For our purposes, it is useful to separate the contribution of the fractional charges to the charging energy from that of the Cooper pairs. We will therefore work in a regime which highlights the role of the former, as done in Ref. 42 for Majorana fermions. If all superconducting islands are connected to a grounded superconductor via a conventional Josephson junction of energy $E_J \gg E_C$, the superconducting phases are pinned to their classical minima, freezing the bosonic degree of freedom associated with Cooper pairs. The charging energy splits the ground state degeneracy by inducing quantum phase slips. In this regime, H_{ch} can be replaced by an effective Hamiltonian of the form⁴³

$$H_\Delta = -\Delta \cos(\pi q + \pi n_{\text{ind}}). \quad (17)$$

The cosine dependence on the charge in this effective Hamiltonian is reminiscent of the Aharonov-Casher effect⁴⁴. When a (Josephson) vortex encircles a superconducting island, it acquires a phase proportional to the charge contained in it. The energy Δ is exponentially small in the ratio E_J/E_C .

Let us now write explicitly the interaction (17) in terms of the PF operators. We denote as $q_{a,k}$ the fractional charge trapped inside the segment of an FTI edge a between the k -th and $(k+1)$ -th ferromagnet. As such, it can be expressed as

$$e^{i\pi q_{a,k}} \equiv P_{\{a,2k\},\{a,2k+1\}} = e^{-i\frac{\pi}{2m}} \alpha_{\{a,2k+1\}}^\dagger \alpha_{\{a,2k\}} \quad (18)$$

In the special case $k = M$, Eq. (18) has to be supplemented with the boundary condition

$$\alpha_{\{a,2M+1\}} = e^{-i\pi q_a} \alpha_{\{a,1\}} \quad (19)$$

where q_a is the total fractional charge along the edge a . Eq. (19) appears naturally in the bosonization description of the PFs as the boundary condition of a closed

edge with total charge q_a surrounding no net magnetic flux,¹⁵ see also Appendix A. This boundary condition constitutes a constraint that the physical states of the system must fulfill, as we will discuss more extensively in the final part of this Section.

In our 2D architecture, each SC gaps out either one or two segments of an FTI edge, depending on whether it lies at the boundary of the system or in the bulk. In the

second case, the total fractional charge q contained in it is the sum of two charges $q_{a,k}, q_{a',k'}$ and can be expressed as $e^{i\pi q} = e^{i\pi(q_{a,k}+q_{a',k'})} = P_{\{a,k\}\{a,k+1\}} P_{\{a',k'\}\{a',k'+1\}}$, since two \mathbb{Z}_{2m} charges operators always commute if they do not share a PF operator. The charging energy takes the form

$$H_{\Delta} = \begin{cases} -\frac{\Delta}{2} \left(e^{i\pi n_{\text{ind}}} e^{-i\pi/2m} \alpha_{\{a,2k+1\}}^{\dagger} \alpha_{\{a,2k\}} + \text{h.c.} \right) & \text{on the boundary} \\ -\frac{\Delta}{2} \left(e^{i\pi n_{\text{ind}}} e^{-i\pi/m} \alpha_{\{a,2k+1\}}^{\dagger} \alpha_{\{a,2k\}} \alpha_{\{a',2k'+1\}}^{\dagger} \alpha_{\{a',2k'\}} + \text{h.c.} \right) & \text{in the bulk} \end{cases} \quad (20)$$

The total charges q_a of the FTI edges may appear in the Hamiltonian (20) as additional phases, due to Eq. (19).

C. Effective Hamiltonian

Adding together the contributions from all islands and junctions, we arrive to an effective Hamiltonian

$$H = \sum_{\text{islands}} H_{\Delta} + \sum_{\text{junctions}} H_J. \quad (21)$$

Each PF of the array belongs to one superconducting island and one junction and therefore it appears twice in the effective Hamiltonian, once in H_{Δ} and once in H_J .

Note that the effective Hamiltonian is the same for the stripe and the tile architectures, which share the same lattice, the same number of parafermions and the same set of local interactions. Nevertheless, the presence of two different sets of commutation rules for the PF operators is enough to give the two systems markedly different properties.

D. Conserved quantities and charge constraints

The two different commutation rules between PFs in the stripe and tile architectures are due to the fact that the Hilbert spaces of the two system are constrained in physically different ways. To see this, notice that the total charge q_a at the edge of each FTI a must be conserved since no term in the Hamiltonian (21) introduces tunneling between different fractional topological insulators. That is,

$$[e^{i\pi q_a}, H] = 0 \quad (22)$$

for every a . Moreover, the total charge of each FTI (edge plus bulk) is not only conserved but also *constrained* to

be an integer multiple of the electron charge e . Thus, if we make the simplifying assumption that there are no fractional excitations trapped in the bulk of the FTIs, we come to the conclusion that all q_a 's must be integer-valued. This requirement restricts the possible eigenvalues of $e^{i\pi q_a}$ to ± 1 , corresponding to the even or odd fermion parity sectors. Without loss of generality, we will assume that each FTI has an even number of electrons, $q_a = 0, \pm 2, \pm 4, \dots$, leading to the set of conditions

$$e^{i\pi q_a} = \prod_{k=1}^M P_{\{a,2k\},\{a,2k+1\}} = 1. \quad (23)$$

This choice amounts to restricting the twisted boundary conditions (19) to the periodic case.

The constraint (23) is violated if a quasiparticle or a quasihole is introduced in the bulk of the FTI. Due to the incompressibility of the FTI liquid, this process is related to the presence of an additional flux quantum $\Phi_0 = h/2e$ piercing the bulk FTI⁴⁵. Thus, we can translate the conservation of electric charge on the edge of the FTI into a conservation of the magnetic flux threaded through the bulk. The latter is measured by the Aharonov-Bohm phase of a quasiparticle performing a counter-clockwise loop along the edge of the FTI. Mathematically, the Aharonov-Bohm phase factor is given by the string product Σ_a of the tunneling operators along such loop,

$$\Sigma_a = \prod_{k=1}^M P_{\{a,2k-1\},\{a,2k\}} \quad (24)$$

which obeys $[\Sigma_a, e^{i\pi q_a}] = 0$, $[\Sigma_a, H] = 0$ and $(\Sigma_a)^{2m} = 1$ and has eigenvalues $\sigma_a = e^{i\pi n/m}$, $n = 0, \dots, 2m-1$. One can derive from Eqs. (3) and (4) the commutation rule

$$\Sigma_a \alpha_{\{a,j\}}^{\dagger} = e^{-i\pi/m} \alpha_{\{a,j\}}^{\dagger} \Sigma_a, \quad (25)$$

which confirms that the operator $\alpha_{\{a,j\}}^{\dagger}$, creating a charge e/m on the edge of the FTI, at the same time adds $-\pi/m$ to the Aharonov-Bohm phase.

Consistently with the constraint (23), we consider as physical only the sector of the full Hilbert space in which

$$\Sigma_a = 1, \quad (26)$$

so that the magnetic flux in each FTI must be a multiple of $4m\Phi_0$. Changing the fluxes that pierce each FTI, we choose a different set of eigenvalues σ_a and select a different physical sector.

The differences between the tile and stripe model originate from the fact that, for fixed system size, the number of FTIs is greater in the tile than in stripe model. This in turns determines the number of independent constraint on the Hilbert space, as reflected in the extent M of the product defining the string operator in Eq. (24) - we recall that $M = 4$ in the tile model and $M = L_x$ in the stripe model. The result is a different dimensionality of the physical sector of the Hilbert space of the two models.

III. MAPPING TO 2D QUANTUM CLOCK MODELS

To highlight the differences between the two models, it helps to use a mapping onto quantum 2D $2m$ -clock models. These models are defined on a 2D lattice where each site \mathbf{r} is a $2m$ -level quantum system and their Hamiltonians possess discrete \mathbb{Z}_{2m} local symmetries. Clock Hamiltonians are defined in terms of degrees of freedom $\sigma_{\mathbf{r}}$ and $\tau_{\mathbf{r}}$ that satisfy

$$\sigma_{\mathbf{r}}^{2m} = \tau_{\mathbf{r}}^{2m} = 1 \quad (27)$$

$$\tau_{\mathbf{r}}^\dagger = \tau_{\mathbf{r}}^{-1}, \quad \sigma_{\mathbf{r}}^\dagger = \sigma_{\mathbf{r}}^{-1}. \quad (28)$$

Operators on any given site have commutation rules similar to those of PFs,

$$\sigma_{\mathbf{r}} \tau_{\mathbf{r}} = e^{i\pi/m} \tau_{\mathbf{r}} \sigma_{\mathbf{r}} \quad (29)$$

but operators on different sites commute. If $m = 1$ these relations are satisfied by Pauli matrices $\sigma_{\mathbf{r}}^z, \sigma_{\mathbf{r}}^x$. In the mathematical literature, the algebra describing PFs is known as a generalized Clifford algebra. Its representation theory has been worked out in detail in Ref. 46, and from it, it is possible to infer a mapping relating the PF operators α and the clock operators σ, τ . This mapping can be achieved via a parafermionic Jordan-Wigner transformation^{26,46,47}.

To each couple of adjacent PFs $\alpha_{\{a,2k\}}, \alpha_{\{a,2k+1\}}$ on the same superconductor, we associate a couple of operators $\sigma_{a,k}, \tau_{a,k}$. These operators therefore live on (a subset of) the links of the square lattice defined by the PFs. The mapping between PFs and clock operators is given by

$$\alpha_{\{a,2k\}} = \kappa_a \sigma_{a,k+1} \prod_{1 \leq l \leq k} \tau_{a,l} \quad (30)$$

$$\alpha_{\{a,2k+1\}} = \kappa_a e^{i\frac{\pi}{2m}} \tau_{a,k+1} \sigma_{a,k+1} \prod_{1 \leq l \leq k} \tau_{a,l} \quad (31)$$




FIG. 3: The transformation defined in Eqs. (30), (31) maps the PF operators living on the sites of a square lattice into a set of clock operators σ, τ defined on the links of the lattice occupied by a superconductor (marked in figure as black dots). Here, as an example, we show the positions of the clock operators in the case of a FTI in the tile model. The mapping between $\alpha_1 \dots, \alpha_8$ and $\sigma_1, \tau_1, \dots, \sigma_4, \tau_4$ shown in this figure is explicitly written down in Eq. (37).

Here κ_a are fractional Klein factors taking care of the commutation rules between parafermions on different edges^{48,49},

$$\kappa_a^{-1} = \kappa_a^\dagger \quad (32)$$

$$\kappa_a \kappa_{a'} = e^{i \text{sgn}(a' - a) \pi / m} \kappa_{a'} \kappa_a. \quad (33)$$

These commutation rules must be compared with Eqs. (3,4,5). Apart from fixing the commutators, the Klein factors do not play a role and drop out from any quadratic operator considered in this paper. The boundary conditions (19) and the constraints (23),(26) are taken into account by setting

$$\sigma_{a,M+1} = \sigma_{a,1} \quad (34)$$

$$\tau_{a,M+1} = \tau_{a,1} \quad (35)$$

$$\prod_{k=1}^M \tau_{a,k} = 1 \quad (36)$$

Let us write down an explicit example of the transformation for the case $M = 4$, relevant for the tile architecture. In this case the relations (30) and (31), dropping the index a and the Klein factors for clarity, read (see also Fig. 3)

$$\begin{aligned} \alpha_8 &= \sigma_1, & \alpha_1 &= e^{i\frac{\pi}{2m}} \tau_1 \sigma_1, \\ \alpha_2 &= \sigma_2 \tau_1, & \alpha_3 &= e^{i\frac{\pi}{2m}} \tau_2 \sigma_2 \tau_1, \\ \alpha_4 &= \sigma_3 \tau_2 \tau_1, & \alpha_5 &= e^{i\frac{\pi}{2m}} \tau_3 \sigma_3 \tau_2 \tau_1, \\ \alpha_6 &= \sigma_4 \tau_3 \tau_2 \tau_1, & \alpha_7 &= e^{i\frac{\pi}{2m}} \tau_4 \sigma_4 \tau_3 \tau_2 \tau_1. \end{aligned} \quad (37)$$

We now rewrite the Hamiltonian in terms of the clock operators. For the Josephson energy, Eq. (11), we obtain

$$H_J = -\frac{J}{2} \left(e^{i\delta/2m} \sigma_{a,k+1}^\dagger \sigma_{a,k} + \text{h.c.} \right), \quad (38)$$




FIG. 4: Layout of the stripe and tile architecture in terms of clock operators σ, τ (black dots), living on the links of the square lattice occupied by a superconductor. Since clock operators at different sites commute, it is not necessary to order the FTI nor to assign an orientation to the FTI edges. However, notice that the clock operators for the two models live on two inequivalent lattices.

while the charging energy, Eq. (20), becomes

$$H_\Delta = \begin{cases} -\frac{\Delta}{2} (e^{i\pi n_{\text{ind}}} \tau_{a,k} + \text{h.c.}) & \text{on the boundary} \\ -\frac{\Delta}{2} (e^{i\pi n_{\text{ind}}} \tau_{a,k} \tau_{a',k'} + \text{h.c.}) & \text{in the bulk} \end{cases} \quad (39)$$

Note that the locality of the interactions is preserved. At this point, it is useful to split the array Hamiltonian of Eq. (21) into bulk and boundary contributions,

$$H = H_{\text{bulk}} + H_{\text{boundary}}, \quad (40)$$

with

$$H_{\text{bulk}} = -\left[\frac{J}{2} \sum_{\text{junctions}} \sigma_{a,k+1}^\dagger \sigma_{a,k} + \frac{\Delta}{2} \sum_{\substack{\text{islands} \\ \in \text{bulk}}} \tau_{a,k} \tau_{a',k'} \right] + \text{h.c.} \quad (41)$$

and

$$H_{\text{boundary}} = -\frac{\Delta}{2} \sum_{\substack{\text{islands} \\ \in \text{bdr}}} (\tau_{a,k} + \tau_{a,k}^\dagger), \quad (42)$$

see also Fig. 4. In writing Eqs. (41),(42) we have, for simplicity, set $n_{\text{ind}} = 0$ for all islands and, in agreement with our choice of the physical sector, $\delta = 0$ for all junctions. Then the couplings J, Δ become purely real and all equal.

Splitting the Hamiltonian into bulk and boundary contributions is useful for studying various boundary conditions. For simplicity, in the remainder of this paper we will set $H_{\text{boundary}} = 0$ and focus on bulk properties of the array, assuming the system size is large enough to justify neglecting H_{boundary} . Also $H_{\text{boundary}} = 0$ corresponds (for any system size) to the exact boundary conditions in case every superconducting island at the boundary of the array is grounded (since in that case $\Delta = 0$ at the boundary).

The Hamiltonian H_{bulk} of Eq. (41) commutes with an extensive set of local operators

$$\xi_s = \sigma_{a,k} \sigma_{a',k'}^\dagger, \quad [\xi_s, H] = 0, \quad (43)$$

associated to every bulk superconducting island s for both the tile and stripe models. Notice however that only those operators ξ_s that commute with the constraints described in the previous section are actual physical symmetries: if $a \neq a'$, the operator (43) moves one fractional charge from one edge to the other, thus violating the charge constraint.

The difference between the tile and the stripe architectures can now be better appreciated, as shown in Fig. 4. The effective Hamiltonian of H_{eff} Eq. (21) associated to either architecture is defined on the square lattice in terms of PFs. In contrast, H_{eff} is defined on inequivalent lattices when represented in terms of clock operators as in Eq. (41). The tile model Hamiltonian H_{tile} is obtained by specializing H_{bulk} to a decorated square lattice, and the stripe model Hamiltonian H_{stripe} is obtained by specializing H_{bulk} to a brickwall lattice instead.

IV. TOPOLOGICAL PHASES AND ORDERS

The quantum phase diagram of the tile and stripe models at zero temperature is controlled by the single parameter J/Δ . In the following we will call *Coulomb-dominated* the regime $\Delta \gg J$ and *Josephson-dominated* the opposite regime $J \gg \Delta$. In this Section we study the two regimes for both models, with a focus on the presence (or absence) of topological order. We dedicate Sec. IV A to the tile model and Sec. IV B to the stripe model. Let us summarize here the main findings.

The Josephson-dominated regime shows no topological features for either model. On one hand, the ground state of the tile model is singly-degenerate in this limit due to the charge constraints. Moreover, exactly at $\Delta = 0$ the ground wave function reduces to a product state of wave functions for local four-body clusters, emphasizing the absence of long-range entanglement. On the other hand, at $\Delta = 0$ the stripe model reduces to a system of decoupled, one-dimensional vector Potts chains in zero transverse field. Hence, in the thermodynamic limit, the

stripe model has ferromagnetic order in the Josephson-dominated regime $J \gg \Delta$. The charge constraints do not suffice to select a unique ground state like for the tile model, but rather correlates the magnetization for pairs of chains.

In the opposite Coulomb-dominated regime, and specifically at $J = 0$, both the tile and stripe model show dramatically increased (relative to $\Delta = 0$) ground-manifold degeneracy. This suggests that at least one phase transition separates the two regimes in the thermodynamic limit for both models.

We will show that the tile model is topologically ordered in the Coulomb-dominated regime, since

1. the degeneracy of its ground manifold depends on the topology of the lattice, and
2. the model has anyonic excitations completely equivalent to those in the qudit toric code with \mathbb{Z}_{2m} discrete symmetry^{39,52}.

The second point is especially noteworthy, since the tile model is akin but not strictly equivalent to the \mathbb{Z}_{2m} toric code by Kitaev³ nor to its generalizations^{39,52}.

Unlike the tile model, that can be defined naturally on a surface of arbitrary genus due to the limited extension of its FTIs, the stripe model fits naturally only open, cylindrical, or periodic (toroidal) boundary conditions. We will see that in the Coulomb-dominated regime its ground state degeneracy is not protected against local operators. Hence we do not consider the stripe model topologically ordered. We will argue nevertheless that the $\Delta \gg J$ regime is characterized by a non-local order parameter, that we will define using a duality mapping the stripe model to the \mathbb{Z}_{2m} lattice gauge theory⁶⁰. Thus, even in the absence of a topological order, the Coulomb-dominated phase of the stripe model can be addressed more generically as a topological phase.

A. A physical realization of \mathbb{Z}_{2m} toric code anyons: the tile model

To analyze the effective Hamiltonian for the tile architecture, it is useful to adopt a decorated square lattice where each FTI sits on a site $\mathbf{r} = i\mathbf{e}_1 + j\mathbf{e}_2$, with (i, j) a pair of integers, see Fig. 5. In this lattice, the array Hamiltonian H_{bulk} of Eq. (41) is given by

$$H_{\text{tile}} = - \left[\frac{J}{2} \sum_{\mathbf{r}} \sum_{i=1}^4 \sigma_{\mathbf{r},i}^\dagger \sigma_{\mathbf{r},i+1} + \frac{\Delta}{2} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} Q_{\langle \mathbf{r}, \mathbf{r}' \rangle} \right] + \text{h.c.} \quad (44)$$

where $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ labels the charging energy terms of the superconducting islands, now sitting on the links of the square lattice between two neighboring diamonds,

$$Q_{\langle \mathbf{r}, \mathbf{r}+\mathbf{e}_1 \rangle} = \tau_{\mathbf{r},3} \tau_{\mathbf{r}+\mathbf{e}_1,1}, \quad Q_{\langle \mathbf{r}, \mathbf{r}+\mathbf{e}_2 \rangle} = \tau_{\mathbf{r},4} \tau_{\mathbf{r}+\mathbf{e}_2,2}. \quad (45)$$




FIG. 5: Panel (a): The notation adopted for the decorated square lattice on which the tile model can be conveniently rearranged. The grey diamonds, sitting on the sites \mathbf{r} of a square lattice, are FTI. To each site \mathbf{r} there correspond four clock operators $\sigma_{\mathbf{r},i}, \tau_{\mathbf{r},i}$, arranged counterclockwise. Blue links are SC, red links are FM. Panel (b): The operator $B_{\mathbf{r}}$ in Eq. (48) is the counter-clockwise product of four $\sigma \sigma^\dagger$ operators around the same plaquette.

Let us now consider the limit $J = 0$ deep in the Coulomb-dominated regime. The system is then in a limit state where tunneling between islands is forbidden. Each superconductor minimizes the charging energy in the space of physical states specified by the charge constraints:

$$Q_{\langle \mathbf{r}, \mathbf{r}+\mathbf{e}_1 \rangle} = Q_{\langle \mathbf{r}, \mathbf{r}+\mathbf{e}_2 \rangle} = 1. \quad (46)$$

These conditions allow for $(2m)$ -fold degeneracy for each superconducting link, corresponding to the presence of the local symmetries (43). However in the sector of physical states we must impose the constraints

$$\prod_{i=1}^4 \tau_{\mathbf{r},i} = 1 \quad (47)$$

derived from Eq. (23). Nearly half of the previous states are then projected out, leaving a ground state manifold of dimension $(2m)^{\#SC/2}$. This number is exact asymptotically in the system size, but depends slightly on the boundary conditions. For example, for periodic boundary conditions the *exact* degeneracy of the ground manifold is $(2m)^{1+(\#SC/2)}$.

The degeneracy of the Coulomb-dominated limit at $J = 0$ is partially lifted when weak tunneling terms are reintroduced, that is, we allow $J \neq 0$. The Coulomb-dominated regime $J \ll \Delta$ can be treated perturbatively by introducing an effective low-energy Hamiltonian affecting only the ground state manifold at $J = 0$. We need to keep only those operators in the perturbative expansion which do not couple the ground state manifold to the excited states. This is a standard technique⁵⁰, and the computation is analogous to the perturbative derivation of the \mathbb{Z}_2 toric code Hamiltonian from Kitaev's honeycomb model⁵¹, so we will only streamline the essential points.

At first order, the perturbation $\sigma_{\mathbf{r},i}^\dagger \sigma_{\mathbf{r},i+1}$ creates two charged $\pm e/m$ excitations on adjacent superconductors,

increasing the energy of the system by an amount $G = 2\Delta(1 - \cos \pi/m)$. Similarly, at all odd orders we obtain terms that we neglect as they do not leave the $J = 0$ ground state manifold invariant. At second order, we obtain only terms describing the tunnelling back and forth of a fractional charge e/m across a single Josephson link. These terms renormalize the ground energy level, that is, they provide an energy offset to the full Hamiltonian. At fourth order we obtain the first relevant contribution. It is a plaquette operator of the form (see Fig. 5)

$$B_{\mathbf{r}} = \left(\sigma_{\mathbf{r},4} \sigma_{\mathbf{r},3}^\dagger \right) \times \left(\sigma_{\mathbf{r}+\mathbf{e}_1,1} \sigma_{\mathbf{r}+\mathbf{e}_1,4}^\dagger \right) \times \left(\sigma_{\mathbf{r}+\mathbf{e}_1+\mathbf{e}_2,2} \sigma_{\mathbf{r}+\mathbf{e}_1+\mathbf{e}_2,1}^\dagger \right) \times \left(\sigma_{\mathbf{r}+\mathbf{e}_2,3} \sigma_{\mathbf{r}+\mathbf{e}_2,2}^\dagger \right), \quad (48)$$

describing the tunnelling of an e/m excitation along a loop of four FTI edges and four superconducting islands. The resulting perturbative Hamiltonian reads

$$H_{\text{pert}}^{\text{tile}} = - \left[\frac{\Delta}{2} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} Q_{\langle \mathbf{r}, \mathbf{r}' \rangle} + \frac{5J^4}{4G^3} \sum_{\mathbf{r}} B_{\mathbf{r}} \right] + \text{h.c.} \quad (49)$$

where we note that, in the case $m = 1$, the coefficient of $B_{\mathbf{r}}$ matches the one obtained in a similar perturbative expansion in Ref. 35, where equivalent plaquette operators are obtained.

Since the operators $B_{\mathbf{r}}$ commute with the charge constraints, the space of physical states for the perturbative Hamiltonian $H_{\text{pert}}^{\text{tile}}$ is left untouched. The bond operators $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ and $B_{\mathbf{r}}$, together with their Hermitian conjugates, constitute a completely commuting set of stabilizers for a qudit surface code³⁹. This surface code protects against every local error that excites a ground state of H_{pert} into a state of higher energy³⁹. In particular, the $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ operators enforce the absence of charge excitations in the superconducting islands, while the $B_{\mathbf{r}}$ operators enforce the absence of flux excitations.

Let us note that the Hamiltonian $H_{\text{pert}}^{\text{tile}}$ is not exactly equivalent to the \mathbb{Z}_{2m} toric code originally discussed by Kitaev in Ref. 3, since the stabilizers $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ and $B_{\mathbf{r}}$ are not projectors. The construction of these operators is instead more closely related to the qudit surface codes introduced in Ref. 39 - although, strictly speaking, $H_{\text{pert}}^{\text{tile}}$ is not equivalent to those models as well, since it is not possible to canonically associate our stabilizers $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ and $B_{\mathbf{r}}$ to vertices and faces of a two-dimensional simplicial complex. Despite these minor differences, however, the topological properties of these models are the same.

The conditions

$$Q_{\langle \mathbf{r}, \mathbf{r}' \rangle} = 1, \quad B_{\mathbf{r}} = 1, \quad \prod_i \tau_{\mathbf{r},i} = 1, \quad (50)$$

specify the ground manifold of the system. Its degeneracy can be determined from symmetry considerations. On the torus (periodic boundary conditions) there are four types of loop symmetries $H_\tau, H_\sigma, V_\tau, V_\sigma$, defined in Fig. 6, associated to non-contractible loops and compatible (commuting) with the charge constraints. It is




FIG. 6: The four non trivial loop operators that define the ground state manifold of the tile model on a torus. The operators H_τ, V_τ are defined as the product of all τ along the path described by the two blue lines, in the order established by the arrows. Similarly, the operators H_σ, V_σ are defined as the product of all $\sigma\sigma^\dagger$ operators along the path given by the red lines. Loop operators corresponding to different paths only differ by a product of stabilizer operators $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ or $B_{\mathbf{r}}$.

interesting to notice (for comparison with other \mathbb{Z}_{2m} surface codes in the literature) that the loop symmetries $H_\tau, V_\tau, (H_\sigma, V_\sigma)$ are disjoint, that is, they do not have any clock degrees of freedom in common. As usual, any two loop symmetries of a given type, τ or σ , associated to equivalent but different non-contractible loops differ only by a product of the stabilizer operators in (50) (or their hermitian conjugate). Hence, in the ground manifold, all these loop symmetries collapse into just four inequivalent ones. These form two non-commuting pairs,

$$H_\tau V_\sigma = e^{-i\pi/m} V_\sigma H_\tau \quad (51)$$

$$H_\sigma V_\tau = e^{-i\pi/m} V_\tau H_\sigma \quad (52)$$

while $[H_\tau, H_\sigma] = [V_\tau, V_\sigma] = [V_\tau, V_\sigma] = [H_\tau, H_\sigma] = 0$. Since $V_\sigma^{2m} = \mathbb{1} = H_\sigma^{2m}$, it follows that each pair identifies $2m$ different ground states, yielding a ground state degeneracy of $(2m)^2$. This is the dimension of the code space defined by $H_{\text{pert}}^{\text{tile}}$.

Our stabilizer code can be adapted to a planar geometry with open boundary conditions along the lines set in Ref. 37. In this case there will be only two non-trivial string operators and thus $2m$ ground states. In planar geometries with g holes, the ground-state manifold degeneracy increases to $(2m)^g$.

The Hamiltonian $H_{\text{pert}}^{\text{tile}}$ has two different types of excitations illustrated in Fig. 7:

1. Two localized charge excitations $\pm e/m$ can be cre-




FIG. 7: Vortex and charge excitations in the tile model, appearing at the ends of open string of τ and $\sigma\sigma^\dagger$ operators respectively. The vortices live on the plaquettes of the lattice, while the charges on the superconducting links. They are mutual Abelian anyons, with an exchange phase $e^{i\pi/2m}$.

ated on two different links by an open string of tunneling operator of the form $S = \prod (\sigma\sigma^\dagger)$. The operator S^\dagger switches the sign of the charges at the end of the open string.

- Two $\pm h/2e$ vortices are created on neighboring plaquettes by one of the two operators τ on the link separating the plaquettes. (The other τ operator belonging to the same link creates the same vortices, but with opposite sign.) The vortices can be moved apart without further energy costs applying a string $T = \prod \tau$ of consecutive τ operators sharing one common plaquette.

Both charge and flux excitations are bosons when considered separately (since different S operators commute with themselves, as well as different T operators). However, when a charge excitation is moved in a loop around a flux excitation, the wavefunction will acquire a (Aharonov-Bohm) phase $e^{i\pi/m}$, implying that charges and flux excitation are mutually Abelian anyons with a fractional exchange phase $e^{i\pi/2m}$. This can be verified by computing the commutator of a pair of S and T strings intersecting each other. Additionally, the underlying \mathbb{Z}_{2m} symmetry allows the presence of multiple excitations of charge ne/m and flux $nh/2e$, with $n = 0, \dots, 2m-1$, created by the n -th power of S and T operators, as in the usual qudit surface codes³⁹.

Let us discuss possible terms that may destroy the topological order. Higher orders in perturbation the-

ory yield larger loop operators which can be decomposed in terms of products of $B_{\mathbf{r}}$ operators and their powers. These higher order terms commute with $H_{\text{pert}}^{\text{tile}}$ and strengthen the absence of fluxes in the plaquettes, leaving the ground state manifold intact. The description breaks down only when the perturbation order L is equal to the system size. At this point, the loop operators H_σ, V_σ are generated in the perturbative expansion, lifting the ground state degeneracy by an energy $O(J^L/\Delta^{L-1})$.

However, we may worry about external perturbations of the form

$$h \sum_{\mathbf{r}, i} (\tau_{\mathbf{r}, i} + \tau_{\mathbf{r}, i}^\dagger), \quad (53)$$

which would break the ground state degeneracy. This perturbation corresponds to an external magnetic field in the vector Potts description and may drive a transition from the topologically ordered phase to a topologically trivial one constituted by a condensate of the vortex excitations. For the \mathbb{Z}_3 toric code, this transition was observed numerically in Ref. 52. A general analysis^{53,54} of the phase diagram of \mathbb{Z}_p ($p = 2, 3, \dots$) Wen-Levin models⁵⁵ suggests that the transition, in the 2+1D transverse-field Potts universality class, is of the first order for any $p > 3$ ($m > 1$), thus easily detectable due to the discontinuity in the energy density. Our model however is of the vector Potts (rather than simple Potts) type and further investigations are required to assert the equivalence of the two cases for generic m .

Finally, let us briefly discuss the Josephson-dominated regime. The topological order in the Coulomb-dominated regime of the tile model disappears when the tunneling terms become comparable to the charging energy. In the opposite extreme limit, $\Delta = 0$, the FTIs decouple and the Hamiltonian is just the sum of the Josephson interactions along each diamond of the lattice in Fig 5. In particular, to minimize the energy, the four clock operators σ for each FTI must be aligned and, considering the charge constraint (36), one obtains that the ground state of each FTI is constituted by an equal superposition of all the polarizations:

$$|\text{GS}\rangle_{\mathbf{r}} = \frac{1}{\sqrt{2m}} \sum_{k=0}^{2m-1} |\sigma_{\mathbf{r},1} = \sigma_{\mathbf{r},2} = \sigma_{\mathbf{r},3} = \sigma_{\mathbf{r},4} = e^{i \frac{k\pi}{m}}\rangle. \quad (54)$$

Thus the total ground state is simply the product of the states $|\text{GS}\rangle_{\mathbf{r}}$ of all the FTIs. Due to the charge constraints, it is unique independently of the topology of the system.

For fixed system size, the ground state will remain non-degenerate also when we consider a small charging energy contribution, $\Delta \ll J$, in the Josephson-dominated regime. In particular the effect of applying all the charging operators $Q_{\langle \mathbf{r}, \mathbf{r}' \rangle}$ in a closed area \mathcal{S} is to orient all the clock operators σ inside \mathcal{S} . The result is the formation a domain wall constituted by all the links along the edge of \mathcal{S} , where the clock operators are not aligned anymore.

The energy cost of the domain wall is proportional to $J\partial\mathcal{S}$, where $\partial\mathcal{S}$ is the number of broken Josephson links along the perimeter of \mathcal{S} . Therefore, the Hamiltonian in the Josephson dominated regime can be seen as the confined phase of a loop model⁵⁶, where the loops are the edges of domains with different spin alignment: J provides a tension to the loops whereas Δ constitutes their kinetic energy. Between the topologically ordered Coulomb-dominated regime and the topologically trivial Josephson-dominated regime other phases may appear and the full phase diagram of the tile model deserves further investigations.

B. The stripe model and the \mathbb{Z}_{2m} Gauge theory

As anticipated at the end of the previous Section, the stripe model is naturally supported on a brick-wall lattice. It is convenient to place the clock degrees of freedom $\sigma_{(i,j)}$ and $\tau_{(i,j)}$ on the sites

$$\{(i,j) \mid i = 0, \dots, L_x - 1, \quad j = 0, \dots, L_y - 1\}$$

of an $L_x \times L_y$ square lattice and distinguish between the two sublattices defined by the conditions $(i+j) = \text{even}$ and $(i+j) = \text{odd}$. In the bulk of this geometry the stripe model becomes the generalization of the XXZ honeycomb compass model³⁶ with \mathbb{Z}_{2m} symmetry:

$$H_{\text{stripe}} = -\left[\frac{\Delta}{2} \sum_{i+j=\text{even}} Q_{(i,j)} + \frac{J}{2} \sum_{j=0}^{L_y-1} \sum_{i=0}^{L_x-2} \sigma_{(i,j)} \sigma_{(i+1,j)}^\dagger\right] + h.c. \quad (55)$$

with

$$Q_{(i,j)} \equiv \tau_{(i,j)} \tau_{(i,j+1)}. \quad (56)$$

Depending on the chosen boundary condition, H_{stripe} must be supplemented with an additional boundary term that we will disregard for the sake of simplicity. The stripe unitary operators

$$S_j S_{j+1} = \prod_{i=0}^{L_x-1} Q_{(i,j)}, \quad j = 0, 2, \dots, L_y - 2 \quad (57)$$

represent the physical constraint on the electric charge of the FTIs, Eq. (36); therefore the physical states $|\Psi\rangle$ must satisfy:

$$S_j S_{j+1} |\Psi\rangle = |\Psi\rangle, \quad j = 0, 2, \dots, L_y - 2. \quad (58)$$

The next task is to specify the physical symmetries of the stripe model. The set of non-trivial unitary operators that commute with H_{stripe} is generated by

$$S_j = \prod_{i=0}^{L_x-1} \tau_{(i,j)}, \quad j = 0, 1, \dots, L_y - 1, \quad (59)$$

$$\xi_{(i,j)} = \sigma_{(i,j)} \sigma_{(i,j+1)}^\dagger, \quad i+j = \text{even}. \quad (60)$$




FIG. 8: Notation adopted to study the stripe model. The black and white sites identify clock operators σ, τ , distinguished by a sublattice degree of freedom. Grey stripes are FTI, blue links are superconductors and red links ferromagnets. Notice that half of the vertical links are missing, thus the stripe model is effectively defined on a brick-wall lattice. The flux threaded through a single plaquette of the lattice is measured by the operator B in the figure (see Eq. (62)). As in the tile model, flux excitations (the oriented blue circles) can be created by an open string of τ operators: however, the geometry constrains their movement in the horizontal direction.

We need to specify those operators in this set that also commute with the charge constraints of Eq. (58). The symmetries S_j trivially satisfy this condition, but they are not all independent in the sector of physical states. We can keep

$$S_j, \quad j = 0, 2, \dots, L_y - 2, \quad (61)$$

as an independent set of one-dimensional symmetries for the stripe model. As explained in the introduction to this section, these symmetries are spontaneously broken in the Josephson-dominated regime (at zero temperature). The effective dimensional reduction displayed by the stripe model in this regime is intimately connected to the one-dimensional symmetries of Eq. (61)⁵⁷.

The local physical symmetries of the stripe model are given by the minimal combination of the operators $\xi_{(i,j)}$ that commute with the charge constraints and they assume the form

$$B_{(i,j)} = \xi_{(i,j)} \xi_{(i+2,j)}^\dagger, \quad i+j = \text{even}. \quad (62)$$

These local symmetries have an immediate interpretation: they describe the Aharonov-Bohm phase associated to the magnetic fluxes threading the plaquettes of the brick-wall lattice (see Fig. 8).

The stripe model has no global symmetries independent of the lower-dimensional symmetries already discussed. The global symmetry $S_1 S_3 \dots S_{L_y-1}$, corresponding to the total charge of the system, is trivially spontaneously broken in the Josephson-dominated regime by the spontaneous breakdown of its one-dimensional constituents. Other global symmetries appear as products of the local symmetries discussed in the previous paragraph, and so cannot be spontaneously

broken either by Elitzur's theorem⁵⁸. This suggests that any ordered phase of the stripe model (outside the the Josephson-dominated limit) must be characterized in terms of a generalized, non-local order parameter⁶¹. However this is not enough to assert that the system shows topological order according to our previous definition based on the topological ground state degeneracy and the presence of anyonic excitation. Rather, in the absence of a Landau local order parameter, we speak more generically of topological phases.

To the purpose of comparing the tile and stripe models in the Coulomb-dominated regime it is useful to perform a perturbative analysis also of the stripe model in the limit $\Delta \gg J$. Just as for the tile model, the first non-trivial term appears at the fourth order in perturbation theory, where the perturbative Hamiltonian of the stripe model becomes:

$$H_{\text{stripe}}^{\text{pert}} = - \sum_{i+j=\text{even}} \left[\frac{\Delta}{2} Q_{(i,j)} + \frac{5J^4}{4G^3} B_{(i,j)} \right] + \text{h.c.} \quad (63)$$

As in the case of the tile model, also for this Hamiltonian it is possible to define localized fractional charge excitations and vortex excitations as end of open strings of τ and $\sigma\sigma^\dagger$ operators. However, for this architecture the vortex excitations can propagate only in the horizontal direction, as can be realized noticing that superconductors of different rows share no common plaquette.

Indeed, in the Hamiltonian $H_{\text{stripe}}^{\text{pert}}$ each row of superconducting islands is decoupled from the others, and is characterized by a ground state degeneracy of $2m$. However, the rows between different FTIs present a non-physical symmetry ξ which does not commute with the constraint (58). Thus, accounting for the charge constraints, the overall degeneracy of the ground states in the physical sector is $(2m)^{\#\text{FTIs}}$. Crucially, this degeneracy is not protected against the local symmetries $\xi_{(i,j)}$. Since these local operators may cause transitions between different ground states, the stripe model does not possess a proper topological order. Despite this fact, the model is characterized by a non-local order parameter, as we will discuss in the following.

To this purpose, and more in general to investigate the phase diagram, it is useful to exploit the bond-algebraic theory of dualities^{31,59} which allows us to study the bulk properties of the constrained stripe Hamiltonian for large system size. Our strategy will be to find a duality (consistent with the constraints), mapping H_{stripe} to a known model. As shown in Refs. 31,59, quantum dualities can be obtained as isomorphisms of bond algebras of interactions preserving locality. In principle, we could study the minimal bond algebra of interactions generated by the bonds $Q_{(i,j)}$ ($i+j = \text{even}$) and $\sigma_{(i,j)}\sigma_{(i+1,j)}^\dagger$ in H_{stripe} . However, a duality derived from this bond algebra, that is, an alternative local representation of these interactions, may not preserve the charge constraints of Eq. (58), because these constraints are not contained in this minimal bond algebra. Hence we consider a larger set of




FIG. 9: The dual lattice on which the \mathbb{Z}_{2m} lattice gauge theory of Eq. (67) is defined. Clock operators now live on those links of the original square lattice which are marked by a blue dot. On this new lattice we find that in perturbation theory the physical interactions are given by the star and plaquette operators B_d and A_d defined in Eqs. (66), (69).

generators

$$Q_{(i,j)}, \quad i = 0, \dots, L_x - 1; \quad j = 0, \dots, L_y - 2; \\ \sigma_{(i,j)}\sigma_{(i+1,j)}^\dagger, \quad i = 0, \dots, L_x - 2; \quad j = 0, \dots, L_y - 1$$

(and Hermitian conjugates) for the stripe model's bond algebra $\mathcal{A}_{\text{stripe}}$. That is, we are including also the bonds $Q_{(i,j)}$ ($i+j = \text{odd}$), which are absent from the Hamiltonian. Such extended bond algebra does contain the charge constraints in Eq. (58); hence a duality for $\mathcal{A}_{\text{stripe}}$ maps these constraints in a well defined fashion either to the identity operator (in which case the duality solves the constraints³¹) or to constraints of the dual model.

The characterization of $\mathcal{A}_{\text{stripe}}$ in terms of relations among its generators reveals the following dual representation of the algebra of interactions,

$$Q_{(i,j)} \xrightarrow{\Phi_d} B_d(i,j), \quad (64)$$

$$\sigma_{(i,j)}\sigma_{(i+1,j)}^\dagger \xrightarrow{\Phi_d} \tau_{(i,j)}, \quad (65)$$

with

$$B_d(i,j) \equiv \begin{cases} \sigma_{(i,j)}^\dagger \sigma_{(i,j+1)}^\dagger & \text{if } i = 0, \\ \sigma_{(i,j)}^\dagger \sigma_{(i,j+1)}^\dagger \sigma_{(i-1,j)} \sigma_{(i-1,j+1)} & \text{otherwise.} \end{cases} \quad (66)$$

Then the dual Hamiltonian, $H_G = \Phi_d(H_{\text{stripe}})$, reads

$$H_G = - \left[\frac{\Delta}{2} \sum_{i+j=\text{even}} B_d(i,j) + \frac{J}{2} \sum_{j=0}^{L_y-1} \sum_{i=0}^{L_x-2} \tau_{(i,j)} \right] + \text{h.c.} \quad (67)$$

and it is unitarily equivalent³¹ to the stripe model.

Up to boundary terms, i.e. the incomplete plaquettes $B_{(0,j)}$ ($j = 0, 2, \dots, L_y - 2$), and a redefinition of the lattice that places the clock degrees of freedom on links rather than sites, we recognize H_G as the Hamiltonian of the \mathbb{Z}_{2m} lattice gauge theory studied in connection to the problem of confinement in QCD⁶⁰ (see Fig. 9). The

local symmetries $B_{(i,j)}$ of the stripe model, Eq. (62), map under duality to

$$B_{(i,j)} \xrightarrow{\Phi_d} A_{d(i,j)}, \quad i+j = \text{even}, \quad (68)$$

with

$$A_{d(i,j)} \equiv \tau_{(i,j)} \tau_{(i+1,j)} \tau_{(i,j+1)}^\dagger \tau_{(i+1,j+1)}^\dagger. \quad (69)$$

As is guaranteed by the formalism, the unitary operators $A_{d(i,j)}$, ($i+j = \text{even}$), commute with the dual Hamiltonian H_G . They correspond to the gauge symmetries of the \mathbb{Z}_{2m} gauge theory and they have the interpretation of measuring the local density of external \mathbb{Z}_{2m} charge. Hence our duality maps the magnetic fluxes in the stripe model, as described by the Aharonov-Bohm operators $B_{(i,j)}$ in Eq. (62), to external \mathbb{Z}_{2m} electric charges in the gauge theory.

At this point we can exploit Eq. (68) to compute the dual representation $H_{\text{stripe}}^{\text{pert } D} = \Phi_d(H_{\text{stripe}}^{\text{pert}})$ of the perturbative Hamiltonian of Eq. (63),

$$H_G^{\text{pert}} = - \sum_{i+j=\text{even}} \left[\frac{\Delta}{2} B_{d(i,j)} + \frac{5J^4}{4G^3} A_{d(i,j)} \right] + \text{h.c.} \quad (70)$$

Remarkably, this is the Hamiltonian for the qudit toric code model^{39,52}. To clarify the notation, notice that due to our definition of the plaquette operator Eq. (66), the two operators $A_{d(i,j)}$, $B_{d(i,j)}$ share one vertical link of the lattice, with $A_{d(i,j)}$ to the right and $B_{d(i,j)}$ to the left of that link (see Fig. 9). This duality, however, is a non-local transformation with respect to the clock operators σ and τ . Thus, even if the spectrum of $H_{\text{stripe}}^{\text{pert}}$ is equivalent to the \mathbb{Z}_{2m} toric code, the stripe model in the Coulomb-dominated regime does not present topological order.

To assert that the phase diagram of the gauge theory and the stripe model are connected by the duality Φ_d , we need to investigate the effect of the duality mapping on the charge constraints of Eq. (58). Remarkably, the charge constraints are *holographic*⁶¹, since they map to boundary constraints for H_G ,

$$S_j S_{j+1} \xrightarrow{\Phi_d} \sigma_{(L_x-1,j)}^\dagger \sigma_{(L_x-1,j+1)}^\dagger, \quad (71)$$

for $j = 0, 2, \dots, L_y - 2$. Then the physical states $|\Psi\rangle_d = \Phi_d|\Psi\rangle$ for H_G , seen as a dual representation of the stripe model, are characterized by the condition

$$\sigma_{(L_x-1,j)}^\dagger \sigma_{(L_x-1,j+1)}^\dagger |\Psi\rangle_d = |\Psi\rangle_d \quad (72)$$

for $j = 0, 2, \dots, L_y - 2$, and not by the standard condition of gauge invariance [that is, invariance under the $A_{(i,j)}$, ($i+j = \text{even}$)].⁶⁰ Despite this difference we will argue in the following that the stripe model and the \mathbb{Z}_{2m} lattice gauge theory share the same phase diagram.

The dual gauge theory H_G presents unusual open boundary conditions. Since the charge constraints are holographic, this is in perspective required to guarantee




FIG. 10: The generalized order parameter (string tension) in both the dual and original lattice.

that the dual charge constraints supported on the boundary commute with the dual Hamiltonian H_G (just as the charge constraints commute with H_{stripe}).

However, the standard view that boundary conditions do not affect the phase diagram in the thermodynamic limit suggests in this case that the dual charge constraints do not affect the phase diagram of H_G , which must then coincide with the standard phase diagram of the \mathbb{Z}_{2m} gauge theory. This view is strengthened by the fact that the dual charge constraints commute with the gauge symmetries of H_G , and so the ground state of H_G will belong to the gauge-invariant sector even after the condition Eq. (72) is imposed. Finally this implies that the stripe model presents the same phase diagram independently on the choice of the charge of each FTI edge in the constraint (23), thus in all the different physical sectors.

In view of these considerations, we can argue that the stripe model shares the phase diagram of the \mathbb{Z}_{2m} gauge theory as described in Ref. 60 (and references therein). It follows that there is indeed one second order phase transition separating the Coulomb-dominated from the Josephson-dominated regime. In the gauge-theory language this transition is understood as a confinement-deconfinement transition. In particular, the Coulomb-dominated regime of the stripe model is dual to the deconfined phase of the gauge theory, while the Josephson-dominated regime corresponds to the confined phase.

The phases of a gauge theory cannot be distinguished by a Landau order parameter⁵⁸. However, for the \mathbb{Z}_{2m} gauge theory dual to the stripe model, there exists a generalized order parameter, the so called *string tension*⁶¹,

that is non-zero in the confining phase and vanishes continuously, but non-analytically at the transition point. The string tension is the expectation value of a string of τ 's in the \mathbb{Z}_{2m} gauge theory, which correspond to an open string of tunneling operators $\sigma\sigma^\dagger$ in the stripe model (see Fig. 10, analogously to the string operators creating charge excitations in the tile model. The ground state expectation value of such string falls continuously, but non-analytically to zero at the second order phase transition separating the Josephson-dominated regime (where it is different from zero) from the Coulomb-dominated regime. On the gauge theory side of the duality, the two phases can be distinguished also by a different scaling of the expectation value of the Wilson loops, which map to sets of τ operators in compact regions in the stripe model.

V. CONCLUSIONS AND OUTLOOK

In summary, we have studied two-dimensional arrays of interacting parafermionic zero modes. Such exotic states form along the edge of fractional topological insulators, at the domain walls between proximity-induced superconducting and ferromagnetic pairing. The dynamic of these zero modes is dictated by two competing effects, the charging energy of each superconducting island and the fractional Josephson tunneling of quasiparticles between different islands.

The underlying fractional edge modes, which are originally described by a helical Luttinger liquid theory, determine crucially the possible lattice geometries and the physical constraints of these parafermionic systems. We have analyzed two possible architectures, the tile and the stripe model. They differ mainly for the fact that in the former the length of the edges is constant, while in the latter it scales with the total size of the architecture. We have discussed how this feature gives rise to different physics, despite the fact that the models are characterized by the same lattice of parafermions and the same local interactions.

The difference is appreciated by exploiting a Jordan-Wigner transformation mapping the parafermionic operators into clock operators. Through this transformation the tile model is described by a Hamiltonian on a decorated square lattice whereas the stripe model becomes a compass model, with \mathbb{Z}_{2m} symmetry, on a brick-wall

lattice.

The tile model presents, at least in perturbation theory, the same topological order of the surface codes characterized by a \mathbb{Z}_{2m} symmetry: it shows the same topological degeneracy of the ground state and the same anyonic excitations. Thus the system we described is a possible physical candidate to the realization of qudit surface code Hamiltonians. It is known that the ground state degeneracy of these systems suffers from thermal fragility^{62,63}. However, we note that the intrinsic noise due to the presence of induced charges on the superconducting islands could help localize the anyonic excitations of the system, and thus to protect the information which may be encoded in the ground states.^{64,65}

The stripe model provides instead a physical realization of the \mathbb{Z}_{2m} lattice gauge theory, a toy model often exploited to study confinement-related problems in lattice field theory. The duality mapping between the stripe model and the lattice gauge theory is non-local in terms of the single clock degrees of freedom, but it is local in terms of the interactions. Unlike in the tile model, a toric code Hamiltonian can only be retrieved in the dual theory, where the operators are non-local. Interestingly enough the physical charge constraints of the FTI edges maps to an holographic constraint in the gauge theory which affects only boundary terms.

To conclude, our work addressed the problem of finding topologically-ordered phases in the phase diagram of these two-dimensional collections of topological defects. Comparing the results obtained for the two architectures, we can see that it is not only the nature of the interactions between the defects (in this case, the \mathbb{Z}_{2m} PF zero modes) that determines the presence of topological order, but also the intrinsic geometry of the topological phases originally generating the defects (in this case, the edges of the fractional topological insulators). Understanding the interplay between this two aspects is crucial to design topologically-ordered architectures.

Acknowledgements.

We would like to thank C.W.J. Beenakker, D. Ferraro, N. Magnoli, G. Ortiz, R. Orus and B. Paredes for insightful discussions. This project was supported by the Dutch Science Foundation NWO/FOM and an ERC Advanced Investigator Grant.

¹ X.-G. Wen, *Quantum Field Theory of Many-Body Systems* (Oxford Univ. Press, Oxford, 2004).

² C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Rev. Mod. Phys.* **80**, 1083 (2008).

³ A. Y. Kitaev, *Ann. Phys.* **303**, 2 (2003).

⁴ A. Y. Kitaev, *Phys. Usp.* **44** (suppl.), 131 (2001).

⁵ G. Volovik, *JETP Lett.* **70**, 609614 (1999).

⁶ N. Read and D. Green, *Phys. Rev. B* **61**, 10267 (2000).

⁷ G. Moore and N. Read, *Nucl. Phys. B* **360**, 362 (1991).

⁸ D. Ivanov, *Phys. Rev. Lett.* **86**, 268 (2001).

⁹ J. Alicea, Y. Oreg, G. Refael, F. von Oppen, and M. P. A. Fisher, *Nature Phys.* **7**, 412 (2011).

¹⁰ J. Alicea, *Rep. Prog. Phys.* **75**, 076501 (2012).

¹¹ C. W. J. Beenakker, *arXiv:1112.1950*.

¹² M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).

- ¹³ X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
- ¹⁴ L. Fu and C. L. Kane, Phys. Rev. B **79**, 161408(R) (2009).
- ¹⁵ N. H. Lindner, E. Berg, G. Refael and A. Stern, Phys. Rev. X **2**, 041002 (2012).
- ¹⁶ D. J. Clarke, J. Alicea and K. Shtengel, Nat. Commun. **4**, 1248 (2013).
- ¹⁷ M. Cheng, Phys. Rev. B **86**, 195126 (2012).
- ¹⁸ A. Vaezi, Phys. Rev. B **87**, 035132 (2013).
- ¹⁹ X. G. Wen, Phys. Rev. B **41**, 12838 (1990).
- ²⁰ D.-H. Lee and X.-G. Wen, Phys. Rev. Lett. **66**, 1765 (1991).
- ²¹ M. Levin and A. Stern, Phys. Rev. Lett. **103**, 196803 (2009).
- ²² M. Barkeshli, C.-M. Jian and X.-L. Qi, Phys. Rev. B **87**, 045130 (2013).
- ²³ Y.-Z. You and X.-G. Wen, Phys. Rev. B **86**, 161107(R) (2012).
- ²⁴ M. Hastings, C. Nayak and Z. Wang, arXiv:1210.5477 (2012).
- ²⁵ Y.-Z. You, C.-M. Jian, and X.-G. Wen, Phys. Rev. B **87**, 045106 (2013).
- ²⁶ P. Fendley, J. Stat. Mech **11**, 11020 (2012).
- ²⁷ See for instance G. Mussardo, *Statistical Field Theory*, Oxford University Press (2009) and references therein.
- ²⁸ E. Fradkin and L. P. Kadanoff, Nucl. Phys. B **170**, 1-15 (1980).
- ²⁹ M. A. Rajapbour and J. Cardy, J. Phys. A **40**, 14703 (2007).
- ³⁰ G. Ortiz, E. Cobanera, and Z. Nussinov, Nucl. Phys. B **854**, 780-814 (2012).
- ³¹ E. Cobanera, G. Ortiz, and Z. Nussinov, Adv. Phys. **60**, 679 (2011).
- ³² A. B. Zamolodchikov and V. Fateev, JETP **62**, 215-225 (1985).
- ³³ G. Delfino, arXiv:1212.3178 (2012).
- ³⁴ C. Xu and L. Fu, Phys. Rev. B **81**, 134435 (2010).
- ³⁵ B. M. Terhal, F. Hassler, and D. P. DiVincenzo, Phys. Rev. Lett. **108**, 260504 (2012).
- ³⁶ Z. Nussinov, G. Ortiz and E. Cobanera, Phys. Rev. B **86**, 085415 (2012).
- ³⁷ S. Bravyi and A. Kitaev, arXiv:quant-ph/9811052 (1998).
- ³⁸ S. Bravyi, B. M. Terhal, and B. Leemhuis, New J. Phys. **12**, 083039 (2010).
- ³⁹ S. S. Bullock and G. K. Brennen, J. Phys. A: Math. Theor. **40** 3481 (2007).
- ⁴⁰ Note that the geometry of the array constrains L_y to be even and L_x to be a multiple of four.
- ⁴¹ L. Fu, Phys. Rev. Lett. **104**, 056402 (2010).
- ⁴² B. van Heck, A. R. Akhmerov, F. Hassler, M. Burrello, and C. W. J. Beenakker, New J. Phys. **14**, 035019 (2012).
- ⁴³ J. Koch, T. M. Yu, J. Gambetta, A. A. Houck, D. I. Schuster, J. Majer, A. Blais, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf, Phys. Rev. A **76**, 042319 (2007).
- ⁴⁴ Y. Aharonov and A. Casher, Phys. Rev. Lett. **53**, 319 (1984).
- ⁴⁵ M. Levin, F. J. Burnell, M. Koch-Janusz, and A. Stern, Phys. Rev. B **84**, 235145 (2011).
- ⁴⁶ T. L. Smith, Pacific J. Math. **149**, 157 (1991) and **149**, 185 (1991).
- ⁴⁷ R. Jagannathan, arXiv:1005.4300 [math-ph] (2010).
- ⁴⁸ R. Guyon, P. Devillard, T. Martin, and I. Safi, Phys. Rev. B **65**, 153304 (2002).
- ⁴⁹ E.-A. Kim, M. J. Lawler, S. Vishveshwara, and E. Fradkin, Phys. Rev. B **74**, 155324 (2006).
- ⁵⁰ S. Bravyi, D. P. DiVincenzo and D. Loss, Ann. Phys. **326**, 2793 (2011).
- ⁵¹ A. Yu. Kitaev, Ann. Phys. (N.Y.) **321**, 2 (2006).
- ⁵² M. D. Schulz, S. Dusuel, R. Orus, J. Vidal, K. P. Schmidt, New J. Phys. **14**, 025005 (2012).
- ⁵³ F. J. Burnell, S. H. Simon and J. K. Slingerland, Phys. Rev. B **84**, 125434 (2011).
- ⁵⁴ F. J. Burnell, S. H. Simon and J. K. Slingerland, New J. Phys. **14**, 015004 (2012).
- ⁵⁵ M. A. Levin and X.-G. Wen, Phys. Rev. B **71**, 045110 (2005).
- ⁵⁶ P. Fendley, Ann. Phys. **323**, 3113 (2008).
- ⁵⁷ Z. Nussinov, G. Ortiz, and E. Cobanera, Ann. Phys. **327**, 2491 (2012).
- ⁵⁸ S. Elitzur, Phys. Rev. D **12**, 3978 (1975).
- ⁵⁹ E. Cobanera, G. Ortiz, and Z. Nussinov, Phys. Rev. Lett. **104**, 020402 (2010).
- ⁶⁰ D. Horn, M. Weinstein, and S. Yankielowicz, Phys. Rev. D **19**, 3715 (1979).
- ⁶¹ E. Cobanera, G. Ortiz, and Z. Nussinov, Phys. Rev. B **87**, 041105(R) (2013).
- ⁶² E. Dennis, A. Kitaev, A. Landahl, and J. Preskill, J. Math. Phys. **43**, 4452 (2002).
- ⁶³ Z. Nussinov and G. Ortiz, Phys. Rev. B **77**, 064302 (2008).
- ⁶⁴ J. R. Wootton and J. K. Pachos, Phys. Rev. Lett. **107**, 030503 (2011).
- ⁶⁵ C. Stark, L. Pollet, A. Imamoglu, and R. Renner, Phys. Rev. Lett. **107**, 030504 (2011).

Appendix A: Description of the system through bosonization

In this Appendix we summarize the main features of the bosonization description of our system and we provide an expression in terms of massless bosonic fields of the parafermion operators α and thus of the related interaction terms. We follow the approach in Refs. 15 and 16, where more details can be found.

In absence of the interactions provided by the superconducting islands and the ferromagnetic insulators, the edge of the FTIs defining our systems, or, equivalently, the double edges of juxtaposed fractional quantum Hall layers with opposite polarization, can be described in terms of the Luttinger liquid Hamiltonian proposed by Wen^{19,20}. In particular the massless edge modes are described by the following Hamiltonian:

$$H_0 = \frac{mv}{2\pi} \int dx \left[(\partial_x \varphi)^2 + (\partial_x \theta)^2 \right] \quad (\text{A1})$$

where v is the speed of the two counterpropagating modes and φ, θ are dual massless bosonic fields obeying the commutation relation:

$$[\varphi(x_1, a, t), \theta(x_2, a', t)] = i \frac{\pi}{m} \delta_{a,a'} \Theta(x_2 - x_1) \quad (\text{A2})$$

where Θ is the Heaviside step function. In particular for each FTI edge a it is possible to define two chiral bosonic fields

$$\varphi_{L/R}(x \pm t, a) \equiv \varphi(x, t, a) \mp \theta(x, t, a) \quad (\text{A3})$$

in such a way that the left and right fermionic modes, with opposite spin polarization, are defined by the operators $\psi_{L/R}(x, t, \mathbf{a}) = \eta_{\mathbf{a}} e^{im\varphi_{L/R}(x \pm vt, \mathbf{a})}$ where $\eta_{\mathbf{a}}$ are fermionic Klein factors. The charge density associated with each edge is $\rho = \partial_x \theta / \pi$, thus, in a closed edge with length \mathcal{L} , the total charge of the edge is related to the boundary conditions of the θ field:

$$\pi q_{\text{tot}}(\mathbf{a}) = \theta(x + \mathcal{L}, \mathbf{a}) - \theta(x, \mathbf{a}) \quad (\text{A4})$$

and analogous conditions relate the field φ with the spin density¹⁵.

For each edge the interaction terms corresponding to the proximity induced superconducting coupling and the backscattering give rise to the interaction Hamiltonian:

$$H_I \propto \int dx [-g_S(x) \cos(2m\varphi) - g_F(x) \cos(2m\theta)] \quad (\text{A5})$$

where g_S and g_F describe respectively the position dependence of the induced superconducting and ferromagnetic couplings.

By selecting a position in the bulk of either a superconducting or a ferromagnetic segment of the edge, if the couplings g are strong enough, one can consider respectively the fields φ and θ as pinned to the semiclassical minima $\varphi_k, \theta_k = 0, \pi/m, 2\pi/m, \dots, (2m-1)\pi/m$. Adopting this approximation and considering the limit of sharp transitions between superconducting and ferromagnetic regions, the parafermion operators can be written as:

$$\alpha_{2k-1, \mathbf{a}} = \kappa_{\mathbf{a}} e^{i(\varphi_{k, \mathbf{a}} - \theta_{k, \mathbf{a}})} \quad (\text{A6})$$

$$\alpha_{2k, \mathbf{a}} = \kappa_{\mathbf{a}} e^{i(\varphi_{k+1, \mathbf{a}} - \theta_{k, \mathbf{a}})} \quad (\text{A7})$$

where $k = 1, \dots, M$ labels the ferromagnets and the superconductors along the edge \mathbf{a} and the tile and stripe models are characterized respectively by $M = 4$ and

$M = L_x$. The fractional Klein factors $\kappa_{\mathbf{a}}$ enforce the correct commutation rules (3) and (4). This definition of the parafermionic modes is not unique (see Refs. 15 and 16 for more detail) but it suffices to our purposes. Finally, for a complete description of the system, it is necessary to take into account the correct boundary conditions.

Through this definition of the parafermionic operators it is easy to verify that the tunneling operators assume the form $e^{-i(\varphi_{k+1, \mathbf{a}} - \varphi_{k, \mathbf{a}})} = P_{\{\mathbf{a}, 2k-1\}, \{\mathbf{a}, 2k\}}$ and thus we recover the usual form for the fractional Josephson interaction (11):

$$H_J = -J \cos \left(\varphi_{k+1, \mathbf{a}} - \varphi_{k, \mathbf{a}} - \frac{\delta}{2m} \right) \quad (\text{A8})$$

Moreover the tunneling string operator $\Sigma_{\mathbf{a}}$ defined in (24) for the two models become $\Sigma_{\mathbf{a}} = \exp(-i(\varphi_{M+1, \mathbf{a}} - \varphi_{1, \mathbf{a}}))$, emphasizing the relation between the boundary conditions of the field φ and the magnetic flux enclosed by the FTI edges. The boundary condition (19) assumes a natural form in the bosonized description due to the boundary relation (A4) since:

$$\alpha_{\{\mathbf{a}, 2M+1\}} = \kappa_{\mathbf{a}} e^{i(\varphi_{M+1, \mathbf{a}} - \theta_{M+1, \mathbf{a}})} = e^{-i\pi q_{\mathbf{a}}} \alpha_{\{\mathbf{a}, 1\}} \Sigma_{\mathbf{a}}^{\dagger}. \quad (\text{A9})$$

Once we apply the parafermionic Jordan-Wigner transformation (30,31) to map the system in a quantum clock model, the previous boundary conditions are translated in the following relations:

$$e^{i\pi q_{\mathbf{a}}} = e^{i(\theta_{M+1, \mathbf{a}} - \theta_{1, \mathbf{a}})} = \prod_{k=1}^M \tau_{k+1}^{\dagger} \quad (\text{A10})$$

$$\Sigma_{\mathbf{a}} = e^{-i(\varphi_{M+1, \mathbf{a}} - \varphi_{1, \mathbf{a}})} = \sigma_{M+1}^{\dagger} \sigma_1 \quad (\text{A11})$$

which generalize the boundary conditions (34,35,36).